

Mycorrhizal infection and high soil phosphorus improve vegetative growth and the female and male functions in tomato

Jennifer L. Poulton¹, David Bryla², Roger T. Koide³, and Andrew G. Stephenson¹

¹Department of Biology and The Intercollege Graduate Degree Program in Ecology, The Pennsylvania State University, University Park, PA 16802 USA;

²USDA-ARS, South Peach Avenue, Fresno, CA 93727 USA; ³Department of Horticulture and The Intercollege Graduate Degree Program in Ecology, 102 Tyson Building, The Pennsylvania State University, University Park, PA 16802 USA

Summary

Author for correspondence:

Roger T. Koide

Tel: +1 814 863 0710

Fax: +1 814 863 6139

Email: rxk13@psu.edu

Received: 28 September 2001

Accepted: 22 November 2001

- To further characterize the effects of mycorrhizal infection and soil phosphorus (P) availability on plant fitness, this study examined their effects on the female and male functions, as well as vegetative growth of tomato (*Lycopersicon esculentum*).
- Two cultivars of tomato were grown in a glasshouse under three treatment combinations: nonmycorrhizal, low P (NMPO); nonmycorrhizal, high P (NMP3); and mycorrhizal, low P (MPO).
- Mycorrhizal infection and high soil P conditions improved several vegetative (leaf area, days until first flower and leaf P concentration) and reproductive traits (total flower production, fruit mass, seed number and pollen production per plant, and mean pollen production per flower). In general, mycorrhizal and P responses were greater for reproductive traits than vegetative traits. In one cultivar, these responses were greater for the male function than the female function.
- Thus, mycorrhizal infection and high soil P conditions enhanced fitness through both the female and male functions. Similar trends were usually observed in the NMP3 and MPO treatments, suggesting that mycorrhizal effects were largely the result of improved P acquisition.

Key words: *Lycopersicon esculentum*, fitness, fruit set, mycorrhiza, pollen, resource allocation, soil fertility, vegetative growth.

© *New Phytologist* (2002) **154**: 255–264

Introduction

Despite the large body of research on arbuscular mycorrhizal (AM) effects on vegetative growth, relatively little attention has been given to its effects on plant reproduction, especially the male function. Because most plants are hermaphrodites (Charnov, 1982), contributing genes to the next generation through both the male and female functions, estimates of reproductive success through the male function are just as important as estimates of success through the female function. Several studies have demonstrated that the male and female functions do not always respond in a similar manner (Horovitz & Harding, 1972; Bertin, 1982). Thus, it cannot be assumed that reproduction will show the same level of response to mycorrhizal infection and high soil P conditions

as vegetative growth, or that the male and female functions of reproduction will respond in similarly.

Because reproductive output is often highly correlated with plant size and nutrient status (Harper & White, 1974; Solbrig, 1981), it is not surprising that mycorrhizal plants usually produce more seeds and fruit than nonmycorrhizal plants (Jensen, 1982, 1983; Schenck & Smith, 1982; Dodd *et al.*, 1983; Koide *et al.*, 1988; Bryla & Koide, 1990a; Carey *et al.*, 1992; Stanley *et al.*, 1993; Vejsadova *et al.*, 1993; Bethlenfalvay *et al.*, 1994, 1997; Lu & Koide, 1994; Subramanian & Charest, 1997). Furthermore, mycorrhizal infection can improve seed quality (seed size and P status) in addition to seed quantity (Koide *et al.*, 1988; Bryla & Koide, 1990a; Lu & Koide, 1991; Koide & Lu, 1992; Lu, 1993; Stanley *et al.*, 1993).

Beyond seed and fruit production, relatively little attention has been given to the effects of mycorrhizal infection on other aspects of plant reproduction. Mycorrhizal infection reduced the time until first flower in tomato (*Lycopersicon esculentum*) and *Abutilon theophrasti* (Bryla & Koide, 1990a; Lu & Koide, 1994). In addition, mycorrhizal infection increased flower bud production in pepper (Dodd *et al.*, 1983) and flower production in soybean (*Glycine max*), tomato, and *A. theophrasti* (Schenk & Smith 1982; Bryla & Koide, 1990a; Lewis & Koide, 1990; Lu & Koide, 1994). Mycorrhizal infection also increased the percentage of flowers producing fruits in tomato and *A. theophrasti* (Bryla & Koide, 1990a; Lu & Koide, 1994).

Since both seeds and pollen depend on the sporophyte for provisioning of nutrients, it is reasonable to expect that mycorrhizal infection and high soil P conditions could improve the male function of reproduction as well as the female function. Past studies have found a positive relationship between environmental factors that affect resource availability (e.g. soil fertility and leaf herbivory) and total pollen production, pollen production per flower, and pollen grain size (Vasek *et al.*, 1987; Devlin, 1989; Young & Stanton, 1990; Stephenson *et al.*, 1992; Lau & Stephenson, 1993, 1994; Stephenson *et al.*, 1994; Quesada *et al.*, 1995; Johannsson & Stephenson, 1998; Stephenson *et al.*, 1998). Furthermore, environmental factors can affect the chemical composition of pollen (Stanley & Linskens, 1974; Baker & Baker, 1979; van Herpen & Linskens, 1981; Lau & Stephenson, 1994; Stephenson *et al.*, 1998). For example, high soil P conditions in the field had an overall positive effect on the male function (staminate flower production, pollen production per flower, pollen grain size, and pollen P concentration) in *Cucurbita pepo* (Lau & Stephenson, 1994). In a preliminary glasshouse study with *C. pepo*, mycorrhizal plants produced marginally more pollen grains per staminate flower and significantly larger pollen grains than nonmycorrhizal plants, with these differences increasing over an 8-wk period (Lau *et al.*, 1995). In another study with *C. pepo* grown in the field, pollen from mycorrhizal plants showed faster *in vitro* pollen tube growth rates than pollen from nonmycorrhizal plants (Stephenson *et al.*, 1998). Recent studies with other species have shown further beneficial effects of mycorrhizal infection on pollen production (Pendleton, 2000; Philip *et al.*, 2001; Poulton *et al.*, 2001b) and pollen performance (Poulton *et al.*, 2001a,b).

In the study reported here, two tomato cultivars were grown in a glasshouse under three treatment combinations: nonmycorrhizal, low P (NMPO); nonmycorrhizal, high P (NMP3); and mycorrhizal, low P (MPO). The effects of mycorrhizal infection and soil P availability were determined for both vegetative (leaf area, days until first flower – a measure of pre-reproductive growth rate – leaf P concentration, and final leaf biomass) and reproductive traits (total flower production per plant), including separate measures of the female (total fruit mass per plant, fruit abortion, fruit P concentration,

mean seed number per fruit and total seed number per plant), and male function (mean pollen production per flower, total pollen production per plant and pollen grain size). Although individual responses to mycorrhizal infection and soil P availability are common in the literature, few studies compare vegetative, female, and male responses within a single study, as described here.

Materials and Methods

In a glasshouse study, two cultivars of tomato (*Lycopersicon esculentum* Mill.), 'VF36' and 'VFNT Cherry', were used to examine the effects of mycorrhizal infection and soil P availability on vegetative growth and the female and male functions of reproduction. Both cultivars were obtained from the Tomato Genetics Research Center at the University of California, Davis, CA, USA. 'VF36' produces two to four flowers per inflorescence and large fruit (mean fruit mass = 6.2 g); 'VFNT Cherry' produces six to eight flowers per inflorescence and small fruit (mean fruit mass = 0.7 g). Both cultivars are self-compatible and show indeterminate growth and reproduction. These cultivars were selected because both have mutations for stem color that can be used to determine paternity in pollen mixture studies (Poulton *et al.*, 2001a).

Soil was collected from a low P (14 ppm Olsen extractable P) field at the Pennsylvania State University Agricultural Experiment Station at Rock Springs, PA, USA. Indigenous mycorrhizal fungi were destroyed by autoclaving the air-dried soil at 105°C for 90 min. Then the soil was stored for 2 wk to avoid the potentially phytotoxic effects of autoclaving (Rovira & Bowen, 1966). In order to improve drainage, the field soil was mixed in a 1 : 3 ratio with sterile medium-grade sand.

On 4 June 1997, 'VF36' and 'VFNT Cherry' tomato seeds were planted in trays containing 50% PGX growing mix (Premier, Riviere-du-Loup, Quebec, Canada), 40% Perlite, and 10% whole-soil inoculum (approx. 75 spores of *Glomus etunicatum* Becker and Gerd. per ml of inoculum) for mycorrhizal seedlings, or 10% autoclaved low P field soil (same soil used to produce inoculum) for nonmycorrhizal seedlings. On 23 June, two seedlings were transplanted into each 'Azalea' pot (3800 cm³; Kord, Bramalea, Ontario, Canada), containing the soil–sand mixture. Mycorrhizal seedlings were inoculated again with 60 ml whole soil inoculum directly around their roots; nonmycorrhizal seedlings received 60 ml autoclaved low P field soil directly around their roots with 5 ml spore washings to obtain comparable nonmycorrhizal microbial inputs (Koide & Li, 1989). Pots were arranged in a randomized block design across three benches. (Two benches had three blocks each; one bench had two blocks.) Soil P treatment was established by watering plants once per week with 500 ml of one-third strength Hoagland's nutrient solution (Machlis & Torrey, 1956) without P for 'low P' plants and with 1 mM KH₂PO₄ for 'high P' plants. A preliminary P

response study had determined the concentration of KH_2PO_4 required for nonmycorrhizal plants to have similar vegetative growth to mycorrhizal plants (Poulton, 2000). This resulted in three treatment combinations: nonmycorrhizal, low P (NMPO); nonmycorrhizal, high P (NMP3); and mycorrhizal, low P (MPO). A drip irrigation system supplied plants with additional water as needed. At 9 wk after transplanting, 400 ml water was supplied to each plant daily (except on the days when Hoagland's solution was provided). Thus, there were 48 pots, with eight replicates per cultivar–treatment combination. This study was performed at the same time as the study in Poulton *et al.* (2001b); some plants were used in both studies.

One randomly selected plant from each pot was removed at 3 wk after transplanting (leaving one plant per pot). At 5 wk after transplanting, a root sample was taken midway between the stem and pot edge in each pot with a no. 15 cork bore. The hole was filled with the autoclaved soil–sand mixture. Roots were rinsed out of the soil samples and initially stored in formaldehyde–acetic acid–ethanol (FAA) solution. Then the roots were cleared and stained with trypan blue to determine level of mycorrhizal infection (NMPO, NMP3 \approx 0%; MPO = 68%) using the modified gridline intersect technique in Koide & Mooney (1987).

The vegetative response to the three treatments was measured as leaf area, days until first flower, leaf P concentration, and final leaf biomass. Starting at 3 wk after transplanting, leaf area measurements were recorded once per week for 3 wk, as described in Bryla & Koide (1990b) and Poulton *et al.* (1998). The first plants began to flower 4 wk after transplanting. For each plant, the date on which its first flower opened was recorded. At 5 wk after transplanting, one fully expanded leaf was collected from each plant for P analysis (the youngest leaf greater than 12 cm long for 'VF36' and 9 cm in long for 'VFNT Cherry'). Leaves were dried at 70°C to constant weight and then ground to obtain subsamples. Weighed subsamples were digested in a H_2SO_4 – H_2O_2 mixture using a Technicon (Tarrytown, NY, USA) block digester (Bryla & Koide, 1990a,b). Then the digested subsamples were analysed for P using the molybdo-phosphate method (Watanabe & Olsen, 1965). Ground leaf tissue from sunflowers was used as reference material to check digestion and analytical procedures with each set of 40 samples. At 18 wk after transplanting, all leaves were harvested from each plant and dried at 70°C to constant weight. Then they were weighed to determine final leaf biomass.


The overall reproductive response to the three treatments was measured as total flower production per plant. In addition, separate measures of the female (total fruit mass per plant, fruit abortion, fruit P concentration, mean seed number per fruit and total seed number per plant) and male function (mean pollen production per flower, total pollen production per plant and pollen grain size) were evaluated. Total flower production per plant was determined by counting

the number of flowers produced per month for the remaining 3 months of the study. All open flowers were vibrated twice per week with a modified electric toothbrush to ensure fruit production via self-pollination. Vibrating a flower at a high frequency causes pollen to be released from the anther cone, saturating the stigmatic surface within the cone. Starting at 6 wk after transplanting, pollen was collected weekly to determine mean pollen production per flower. On each plant, all open flowers were vibrated, and pollen for that plant was collected in a single gelatin caplet. Pollen samples were dried at 40°C and stored in 20-ml vials for future analysis. Individual pollen samples were then weighed on a scale with precision to 0.001 mg. Mean pollen production per flower was calculated by dividing mass of the pollen sample by the number of flowers vibrated on that plant. Pollen counts performed on microscope slides showed a strong correlation ($R^2 = 0.97$, $n = 36$) with mass (Poulton, 2000). Total pollen production per plant was calculated as number of flowers \times mean pollen production per flower. Pollen samples were collected monthly to measure pollen grain size. One pollen load was transferred on the flat end of a wire to a microscope slide and was stained with acetocarmine (Kearns & Inouye, 1993). On each slide, the diameters of the first 20 pollen grains encountered on a grid were measured with a micrometer. As fruit ripened, they were harvested and dried at 70°C to constant weight. Dried fruit were weighed to determine total fruit mass per plant. Fruit abortion was calculated as ((number of flowers – number of mature fruit)/number of flowers) \times 100. The number of seeds per fruit was recorded from a representative sample of the fruit (30% of the fruit produced by each cultivar–treatment combination). Total seed production per plant was calculated as number of fruit \times mean seed number per fruit. Fruit P concentration was determined as previously described for leaf tissue.

For most vegetative and reproductive traits, fixed effects analyses of variance (general linear model; Minitab, 1997) were performed with two factors and their interaction: cultivar (two levels) and treatment (three levels). Time (three levels) was also included in the analysis of variance for leaf area. Least square means were calculated whenever sample size varied among cultivar–treatment combinations. Bonferroni pairwise comparisons were used to detect significant differences among the treatments within a cultivar. In addition, multivariate analyses of variance (General MANOVA; Minitab, 1997) were performed for vegetative traits, measures of the female function and measures of the male function, in order to detect overall significant treatment effects. Total flower production per plant was included in both MANOVAs for reproductive traits. Mycorrhizal response was calculated as ((MPO – NMPO mean)/NMPO mean) \times 100 (Bryla & Koide, 1990b). Similarly, P response was calculated as ((NMP3 – NMPO mean)/NMPO mean) \times 100. Leaf area (at 5 wk), total flower production per plant, total seed production per plant and total pollen production per plant were used to

Table 1 Analysis of variance of leaf area (measured at weeks 3, 4 and 5) in 'VF36' and 'VFNT Cherry'


Source	df	F	P
Time	2	184.71	0.001
Cultivar	1	1763.77	0.001
Treatment	2	176.83	0.001
Time × cultivar	2	50.99	0.001
Time × treatment	4	5.87	0.001
Cultivar × treatment	2	19.21	0.001
Time × cultivar × treatment	4	0.48	0.749

**Fig. 1** Leaf area (measured at weeks 3, 4, and 5) in (a) 'VF36' and (b) 'VFNT Cherry'. Means (\pm SE) are given. Solid line, NMPO (nonmycorrhizal, low P); large dashes, NMP3 (nonmycorrhizal, high P); small dashes, MPO (mycorrhizal, low P).

evaluate plant response to mycorrhizal infection and high soil P conditions.

Results

As expected, cultivar had a significant effect on most vegetative and reproductive traits throughout this study. In

**Fig. 2** Relationship between leaf area (cm²) and leaf P concentration at week 5 in (a) 'VF36' and (b) 'VFNT Cherry'. Circles, NMPO (nonmycorrhizal, low P); squares, NMP3 (nonmycorrhizal, high P); triangle, MPO (mycorrhizal, low P).

both cultivars, NMP3 and MPO plants had greater leaf area than NMPO plants, with these differences among treatments increasing over time (Table 1, Fig. 1). Treatment also had a significant effect on days until first flower (Table 2). In the cv. VF36, NMP3 and MPO plants began to flower significantly faster than NMPO plants. In the cv. VFNT Cherry, NMP3 plants began to flower significantly faster than NMPO plants (MPO plants were intermediate). Treatment also had a significant effect on leaf P concentration and final leaf biomass (Table 2). In both cultivars, leaves from NMP3 plants had a significantly higher P concentration than leaves from NMPO plants (MPO plants were intermediate). In addition, NMP3 plants had greater final leaf biomass than NMPO and MPO plants. In both cultivars, the relationship between leaf area and leaf P concentration at week 5 varied among the treatments, although there was always a positive correlation ('VF36' $R^2 = 0.55$; 'VFNT Cherry' $R^2 = 0.66$; Fig. 2). In the multivariate analysis of variance, there was an overall significant treatment effect for vegetative traits, especially in 'VFNT Cherry' (MANOVA, $F = 18.88$, $P = 0.001$).

Table 2 Vegetative traits in 'VF36' and 'VFNT Cherry'. Days until first flower, leaf P concentration (at week 5), and final leaf biomass (g) are given as means and SE (in parentheses)

Cultivar	Treatment	Days until first flower	Leaf P concentration	Final leaf biomass
VF36	NMPO	47.1 (2.6)a	0.17 (0.03)a	4.9 (0.8)a
	NMP3	37.4 (0.5)b	0.27 (0.02)b	8.2 (0.7)b
	MPO	40.9 (0.8)b	0.21 (0.01)ab	5.5 (0.4)a
VFNT Cherry	NMPO	37.0 (2.9)a	0.17 (0.02)a	3.0 (0.3)a
	NMP3	29.0 (0.6)b	0.31 (0.03)b	3.7 (0.3)b
	MPO	34.9 (1.1)ab	0.25 (0.03)ab	2.5 (0.2)a
ANOVA <i>P</i> -values	Cultivar	0.001	0.108	0.001
	Treatment	0.001	0.001	0.001
	Cultivar × treatment	0.489	0.541	0.052

Different letters within a cultivar indicate a significant difference at $P = 0.05$. NMPO, nonmycorrhizal, low P; NMP3, nonmycorrhizal, high P; MPO, mycorrhizal, low P.

Table 3 Measures of the female function of reproduction in the 'VF36' and 'VFNT Cherry' cultivars. Total flower production per plant, total fruit mass per plant (g), fruit abortion and fruit P concentration are given as means and SE (in parentheses); mean seed number per fruit and total seed number per plant are given as least square means and SE (in parentheses)

Cultivar	Treatment	Total flower production per plant	Total fruit mass per plant	Fruit abortion
VF36	NMPO	7.9 (1.5)a	9.1 (1.4)a	75.9 (5.8)a
	NMP3	12.8 (1.5)b	18.6 (1.3)b	71.1 (3.3)a
	MPO	12.5 (1.7)b	13.6 (1.3)ab	81.0 (4.5)a
VFNT Cherry	NMPO	29.0 (3.3)a	5.2 (1.3)a	68.9 (6.8)a
	NMP3	61.3 (3.8)c	16.8 (1.2)b	59.8 (1.9)a
	MPO	42.3 (3.7)b	13.9 (0.5)b	53.3 (4.8)a
ANOVA <i>P</i> -values	Cultivar	0.001	0.078	0.001
	Treatment	0.001	0.001	0.320
	Cultivar × treatment	0.001	0.249	0.084

		Fruit P concentration	Mean seed number per fruit	Total seed number per plant
VF36	NMPO	0.22 (0.02)a	89.8 (18.0)a	130.2 (45.0)a
	NMP3	0.38 (0.02)b	75.8 (11.6)a	273.1 (35.6)b
	MPO	0.25 (0.03)a	89.4 (15.2)a	195.0 (45.0)ab
VFNT Cherry	NMPO	0.17 (0.02)a	38.4 (4.8)a	392.6 (110.0)a
	NMP3	0.29 (0.01)b	54.3 (2.8)b	1300.9 (102.9)c
	MPO	0.22 (0.02)a	48.0 (2.9)ab	879.4 (102.9)b
ANOVA <i>P</i> -values	Cultivar	0.001	0.001	0.001
	Treatment	0.001	0.805	0.001
	Cultivar × treatment	0.212	0.091	0.001

Different letters within a cultivar indicate a significant difference at $P = 0.05$. NMPO, nonmycorrhizal, low P; NMP3, nonmycorrhizal, high P; MPO, mycorrhizal, low P.

Treatment had a significant effect on total flower production per plant, especially in 'VFNT Cherry' (Table 3). In the 'VF36', NMP3 and MPO plants produced significantly more flowers than NMPO plants. In 'VFNT Cherry', NMP3 plants produced significantly more flowers than MPO plants, which in turn produced significantly more flowers than NMPO plants (i.e. NMP3 > MPO > NMPO). Treatment also had a significant effect on total fruit mass per plant (Table 3). In 'VF36', NMP3 plants had significantly greater total fruit mass than NMPO plants (MPO plants were intermediate). In 'VFNT Cherry', NMP3 and MPO plants

had significantly greater total fruit mass than NMPO plants. Treatment did not have a significant effect on fruit abortion (Table 3). However, there was a significant treatment effect on fruit P concentration (Table 3). In both cultivars, fruit from NMP3 plants had a significantly higher P concentration than fruit from NMPO and MPO plants. Treatment also had a significant effect on total seed number per plant, especially in 'VFNT Cherry' (Table 3). In 'VF36', NMP3 plants produced significantly more seeds than NMPO plants (MPO plants were intermediate). In 'VFNT Cherry', NMP3 plants produced significantly more seeds than MPO plants, which in

Table 4 Measures of the male function of reproduction in the VF36 and VFNT Cherry cultivars. Mean pollen production per flower (mg), total pollen production per plant (mg), and pollen grain size (microns) are given as least square means and SE (in parentheses)

Cultivar	Treatment	Mean pollen production per flower	Total pollen production per plant	Pollen grain size
VF36	NMPO	0.132 (0.063)a	1.26 (0.81)a	28.3 (0.4)a
	NMP3	0.329 (0.063)b	4.75 (0.89)b	28.3 (0.4)a
	MPO	0.284 (0.055)b	3.63 (0.70)b	27.7 (0.4)a
VFNT Cherry	NMPO	0.025 (0.009)a	0.86 (0.43)a	25.4 (0.3)a
	NMP3	0.039 (0.006)ab	2.41 (0.34)b	25.7 (0.2)a
	MPO	0.054 (0.007)b	2.26 (0.36)b	25.8 (0.2)a
ANOVA <i>P</i> -values	Cultivar	0.001	0.010	0.001
	Treatment	0.022	0.001	0.675
	Cultivar × treatment	0.071	0.328	0.283

Different letters within a cultivar indicate a significant difference at $P = 0.05$. NMPO, nonmycorrhizal, low P; NMP3, nonmycorrhizal, high P; MPO, mycorrhizal, low P.

Table 5 Mycorrhizal and P responses (%) of selected vegetative and reproductive traits in 'VF36' and 'VFNT Cherry'. Least square means and (SE) of responses and *P*-values from analyses of variance are given

Trait	'VF36'			'VFNT Cherry'				
	Mycorrhizal response	<i>P</i>	P response	Mycorrhizal response	<i>P</i>	P response	<i>P</i>	
Leaf area (at 5 weeks)	50.5 (4.6)	0.001	55.4 (2.6)	0.001	88.6 (8.0)	0.001	95.3 (8.3)	0.001
Total flower production per plant	58.2 (20.9)	0.013	61.4 (18.9)	0.007	45.7 (12.6)	0.004	111.2 (13.2)	0.001
Total seed production per plant	49.8 (32.9)	0.100	109.8 (31.7)	0.005	123.8 (13.9)	0.001	231.0 (34.0)	0.001
Total pollen production per plant	188.1 (49.7)	0.003	277.0 (107.0)	0.031	162.5 (60.3)	0.018	180.4 (26.5)	0.001

turn produced significantly more seeds than NMPO plants (i.e. NMP3 > MPO > NMPO). In addition, NMP3 plants produced significantly more seeds per fruit than NMPO plants (MPO plants were intermediate) in this cultivar (Table 3). In the multivariate analysis of variance, there was an overall significant treatment effect for measures of the female function, especially in 'VFNT Cherry' (MANOVA, $F = 8.83$, $P = 0.001$).

Treatment had a significant effect on mean pollen production per flower (Table 4). In 'VF36', NMP3 and MPO plants produced significantly more pollen per flower than NMPO plants. In 'VFNT Cherry', MPO plants produced significantly more pollen per flower than NMPO plants (NMP3 plants were intermediate). For total pollen production per plant, differences among the treatments were even more significant (Table 4). In both cultivars, NMP3 and MPO plants produced significantly more pollen per plant than NMPO plants. However, treatment did not have a significant effect on pollen grain size (Table 4). In the multivariate analysis of variance, there was an overall significant treatment effect for measures of the male function, especially in 'VF36' (MANOVA $F = 3.02$, $P = 0.015$).

Mycorrhizal and P responses (%) of selected vegetative and reproductive traits (Table 5) were calculated in order to compare levels of response across factors (e.g. mycorrhizal infection vs high soil P conditions, vegetative traits vs reproductive

traits, female vs male, 'VF36' vs 'VFNT Cherry'). In both cultivars, P responses were higher than mycorrhizal responses for all vegetative and reproductive traits. All mycorrhizal and P responses were significant, except for the mycorrhizal response for total seed production per plant in 'VF36'. In general, reproductive responses to mycorrhizal infection and high soil P conditions were greater than vegetative responses. In 'VF36', mycorrhizal and P responses for total pollen production per plant were significantly greater than responses for leaf area and total seed production per plant (mycorrhizal response ANOVA, $F = 5.26$, $P = 0.016$; P response ANOVA, $F = 5.02$, $P = 0.021$). In 'VFNT Cherry', there was no significant difference among mycorrhizal responses (ANOVA, $F = 1.22$, $P = 0.316$). However, P responses for total seed production per plant and total pollen production per plant were significantly greater than the P response for leaf area (ANOVA, $F = 7.04$, $P = 0.005$).

Discussion

In addition to the anticipated effects on vegetative growth, this study clearly demonstrates that mycorrhizal infection and high soil P conditions also improve the female and male functions of reproduction. In many species, mycorrhizal infection enhances P uptake from the soil, thereby improving vegetative growth (Smith & Read, 1997). In this study,

similar trends in NMP3 and MPO plants and the positive correlation between leaf area and leaf P concentration across the treatments suggest that mycorrhizal effects on vegetative growth (and ultimately reproduction) were largely the result of improved P acquisition. Similarly, in separate studies with *C. pepo*, mycorrhizal infection and high soil P conditions had similar beneficial effects on the male function (Lau & Stephenson, 1994; Lau *et al.*, 1995). However, in *Cucurbita foetidissima*, low-P + AMF (arbuscular–mycorrhizal fungi) plants produced significantly more staminate flowers than low-P and high-P plants (Pendleton, 2000).

In this study, mycorrhizal infection and high soil P conditions increased both leaf area (measured before flowering) and total flower production per plant. In many species, flower production tends to increase with plant size (Jackson & Sweet, 1972; Willson *et al.*, 1979; Lloyd, 1980; Aker, 1982; Lee & Bazzaz, 1982). Similar mycorrhizal effects on total flower production have been reported for other tomato accessions (Bryla & Koide, 1990a) as well as other species (Schenck & Smith, 1982; Dodd *et al.*, 1983; Koide *et al.*, 1988; Lewis & Koide, 1990; Lu & Koide, 1994; Ganade & Brown, 1997). However, mycorrhizal effects on flower production independent of plant size have been observed in other studies. For example, Pendleton (2000) found that mycorrhizal infection increased staminate flower production in *C. foetidissima* with no effect on plant biomass.

Mycorrhizal infection and high soil P conditions improved fitness through the female function in this study. These treatments increased total fruit mass per plant by increasing total flower production per plant. Reproductive output is often highly correlated with plant size and nutrient status (Harper & White, 1974; Solbrig, 1981) and total flower production (Stephenson, 1992). Furthermore, mycorrhizal infection and high soil P conditions increased total seed production per plant by increasing fruit production (in both cultivars) and seed number per fruit (in one cultivar). In other tomato accessions, mycorrhizal infection also increased fruit and seed production and increased the percentage of flowers producing fruit (Bryla & Koide, 1990a). Similar mycorrhizal effects on fruit and seed production have been observed in several species (see Smith & Read, 1997).

In this study, mycorrhizal infection and high soil P conditions also improved fitness through the male function. These treatments increased total pollen production per plant by increasing both total flower production per plant and pollen production per flower. In natural populations, siring success is often positively correlated with flower and pollen production (Schoen & Steward, 1986; Devlin *et al.*, 1992). Moreover, pollinator attraction and subsequent pollen dissemination tend to increase with the size of the floral display (Schaffer & Schaffer, 1979; Stephenson, 1979; Willson *et al.*, 1979; Schemske, 1980a,b; Davis, 1981; Patton & Ford, 1983). Other studies have also found that mycorrhizal infection and high soil P conditions improve the male function by

increasing both flower and pollen production (Lau & Stephenson, 1994; Lau *et al.*, 1995; Stephenson *et al.*, 1998; Poulton *et al.*, 2001b). For example, Pendleton (2000) found that mycorrhizal infection increased staminate flower production, and thus pollen production, in *C. foetidissima*. However, Philip *et al.* (2001) found that mycorrhizal infection increased pollen production per anther and per flower in *Lythrum salicaria* without affecting flower production.

In this study, mycorrhizal infection and high soil P conditions also decreased the number of days until first flower (related to plant size), as seen in other tomato accessions (Bryla & Koide, 1990a). In natural populations of outcrossing plants, earlier flowering could potentially improve fitness through both the female and male functions. In species with indeterminate reproduction, earlier flowering can increase total flower production by lengthening the reproductive period. In addition, plants that begin to flower earlier also begin to develop fruit earlier, possibly pre-empting resources from neighboring plants. Moreover, the pollen grains produced by the first flowers in a population have the first opportunity to fertilize ovules at a time when developing fruit are most likely to reach maturity.

Although not measured in this study, mycorrhizal infection and high soil P conditions also increase fitness through the female and male functions by improving the quality of seeds and pollen produced, as well as the quantity. During seed and pollen development, environmental conditions that affect resource availability to the sporophyte (e.g. soil fertility and leaf herbivory) can influence the quantity and quality of seeds (Roach & Wulff, 1987) and pollen produced (Stephenson *et al.*, 1994; Delph *et al.*, 1997). Several studies have found that mycorrhizal infection improves seed quality (seed size and P status), thus enhancing offspring vigor (Lewis & Koide, 1990; Koide & Lu, 1992; Lu & Koide, 1994; Shumway & Koide, 1994a,b; Heppel *et al.*, 1998). In addition, other studies have found that mycorrhizal infection and high soil P conditions improve pollen quality (*in vitro* pollen tube growth rates), thus enhancing siring success in both pollen mixture studies and experimental arrays (Lau & Stephenson, 1994; Stephenson *et al.*, 1998; Poulton *et al.*, 2001a,b).

The P responses in this study were higher than mycorrhizal responses for all vegetative and reproductive traits in Table 5. This result simply indicates that the level of P selected for NMP3 plants (based on an earlier P response study, Poulton, 2000) may have been too high. In addition, NMP3 plants received nutrient solution containing additional P throughout the growing season, while mycorrhizal hyphae were probably less effective at increasing P uptake in MPO plants as their roots became pot-bound at the end of the growing season. This reduction in the effectiveness of mycorrhizal hyphae occurred during the portion of the growing season when fruit development was at its highest, thus affecting total seed production per plant. However, the NMP3 and MPO treatments were not significantly different for most traits.

In both cultivars, mycorrhizal and P responses for reproductive traits were generally greater than responses for leaf area. However, the cultivars differed in their responses for the female and male functions. In the VF36 cultivar, mycorrhizal and P responses for total pollen production per plant were significantly greater than responses for total seed production per plant. Similarly, Pendleton (2000) found that mycorrhizal infection greatly increased staminate flower production in *C. foetidissima* with no effect on pistillate flower production. Acquisition of P (via mycorrhizal infection or other means) may be especially important for pollen grains, which showed the highest P requirement of all reproductive tissues in *Sidalcea oregana* (Ashman & Baker, 1992). For example, stored phytate in mature pollen grains is hydrolysed into phosphate and myoinositol, which are used by the pollen tube for cell wall and membrane synthesis (Jackson & Linskens, 1982; Dickenson & Lin, 1986). However, mycorrhizal and P responses of the male and female functions were not significantly different in 'VFNT Cherry'.

Despite differences in plant morphology, flowers per inflorescence, and fruit size, 'VF36' and 'VFNT Cherry' showed similar responses to treatment for many vegetative and reproductive traits measured in this study. Other tomato accessions have demonstrated significant variation in response to mycorrhizal infection (but no negative response, Bryla & Koide, 1990b). In general, cultivated varieties of tomato were more responsive to mycorrhizal infection than wild accessions, which may be better adapted to poor soil fertility (Koide *et al.*, 1988).

The evolutionary significance of mycorrhizal infection lies in its effects on plant reproduction. Selection for a trait, such as being mycorrhizal, can occur if that trait positively influences fitness. As demonstrated in the preceding study, mycorrhizal infection, similar to high soil P conditions, can increase fitness through both the female and male functions. The fact that most terrestrial angiosperm species are mycorrhizal (Law, 1988) may indicate that mycorrhizal infection does indeed increase the fitness of host plants in natural populations (Koide, 1998). Furthermore, nonmycotrophic species have developed morphological and physiological mechanisms for increasing P acquisition (e.g. proteoid roots, high root and root hair density, and acidification of the rhizosphere; Koide, 1998). Thus, the need to acquire P from the soil (by mycorrhizal infection or other means) has influenced the evolution of plant species. Understanding the impact of mycorrhizal infection on all aspects of plant reproduction, as well as vegetative growth, will become increasingly important as research on the use of mycorrhizas in agriculture, forestry, and land reclamation continues.

Acknowledgements

We thank The Department of Horticulture for the use of their glasshouses and R. Oberheim and his staff at The

Pennsylvania State University Agricultural Experiment Station at Rock Springs, PA, USA, the Tomato Genetic Resources Center, University of California, Davis, CA, USA, for supplying the tomato seeds and R. Smith, C. Di Folco and C. Chuckalovcak for field, glasshouse and laboratory assistance. This research was supported by NSF grant IBN #9419722 to A.G.S. and R.T.K., and Hill–Hill Fellowships to J.L.P.

References

- Aker CL. 1982. Regulation of flower, fruit and seed production by a monocarpic perennial, *Yucca whipplei*. *Journal of Ecology* **70**: 357–372.
- Ashman T-L, Baker I. 1992. Variation in floral sex allocation with time of season and currency. *Ecology* **73**: 1237–1243.
- Baker HG, Baker I. 1979. Starch in angiosperm pollen grains and its evolutionary significance. *American Journal of Botany* **66**: 591–600.
- Bertin RI. 1982. Paternity and fruit production in trumpet creeper. *American Naturalist* **119**: 694–709.
- Bethlenfalvay GJ, Mihara KL, Schreiner RP. 1994. Mycorrhizae alter protein and lipid contents and yield of pea seeds. *Crop Science* **34**: 998–1003.
- Bethlenfalvay GJ, Schreiner RP, Mihara KL. 1997. Mycorrhizal fungi effects on nutrient composition and yield of soybean seeds. *Journal of Plant Nutrition* **20**: 581–591.
- Bryla DR, Koide RT. 1990a. Regulation of reproduction in wild and cultivated *Lycopersicon esculentum* Mill. by vesicular–arbuscular mycorrhizal infection. *Oecologia* **84**: 74–81.
- Bryla DR, Koide RT. 1990b. Role of mycorrhizal infection in the growth and reproduction of wild vs cultivated plants. II. Eight wild accessions and two cultivars of *Lycopersicon esculentum* Mill. *Oecologia* **84**: 82–92.
- Carey PD, Fitter AH, Watkinson AR. 1992. A field study using the fungicide benomyl to investigate the effect of mycorrhizal fungi on plant fitness. *Oecologia* **90**: 550–555.
- Charnov E. 1982. *Theory of sex allocation*. Princeton, NJ, USA: Princeton University Press.
- Davis MA. 1981. The effect of pollinators, predators and energy constraints on the floral ecology and evolution of *Trillium erectum*. *Oecologia* **48**: 400–406.
- Delph LF, Johannsson MH, Stephenson AG. 1997. How environmental factors affect pollen performance: ecological and evolutionary perspectives. *Ecology* **78**: 1632–1639.
- Devlin B. 1989. Components of seed and pollen yield of *Lobelia cardinalis*: variation and correlations. *American Journal of Botany* **76**: 204–214.
- Devlin B, Clegg J, Ellstrand NC. 1992. The effect of flower production on male reproductive success in wild radish populations. *Evolution* **46**: 1030–1042.
- Dickenson DB, Lin JJ. 1986. Phytases of germinating lily pollen. In: Mulcahy DL, Mulcahy GB, Ottaviano E, eds. *Biotechnology and ecology of pollen*. New York, NY, USA: Springer, 357–352.
- Dodd J, Krikun J, Haas J. 1983. Relative effectiveness of indigenous populations of vesicular–arbuscular mycorrhizal fungi from four sites in Negev, Israel. *Israel Journal of Botany* **32**: 10–21.
- Ganade G, Brown VK. 1997. Effects of below-ground insects, mycorrhizal fungi and soil fertility on the establishment of *Vicia* in grassland communities. *Oecologia* **109**: 374–381.
- Harper JL, White J. 1974. The demography of plants. *Annual Review of Ecology and Systematics* **5**: 419–463.
- Heppel KB, Shumway DL, Koide RT. 1998. The effect of mycorrhizal infection of *Abutilon theophrasti* on competitiveness of offspring. *Functional Ecology* **12**: 171–175.
- van Herpen MMA, Linskens HF. 1981. Effect of season, plant age, and temperature during plant growth on compatible and incompatible pollen tube growth in *Petunia hybrida*. *Acta Botanica Neerlandica* **30**: 209–218.

- Horovitz A, Harding J. 1972. The concept of male outcrossing in hermaphrodite plants. *Heredity* 29: 223–236.
- Jackson JF, Linskens HF. 1982. Phytic acid in *Petunia hybrida* pollen is hydrolysed during germination by a phytase. *Acta Botanica Neerlandica* 31: 441–447.
- Jackson DI, Sweet GB. 1972. Flower initiation in temperate woody plants. *Horticulture Abstracts* 49: 9–24.
- Jensen A. 1982. Influence of four vesicular–arbuscular mycorrhizal fungi on nutrient uptake and growth in barley (*Hordeum vulgare*). *New Phytologist* 90: 45–50.
- Jensen A. 1983. The effect of indigenous vesicular–arbuscular mycorrhizal fungi on nutrient uptake and growth of barley in two Danish soils. *Plant Soil* 70: 155–163.
- Johannsson MH, Stephenson AG. 1998. Effects of temperature during microsporogenesis on pollen performance in *Cucurbita pepo* L. (Cucurbitaceae). *International Journal of Plant Science* 159: 616–626.
- Kearns CA, Inouye DW. 1993. *Techniques for pollination biologists*. Niwot, CO, USA: University Press of Colorado.
- Koide RT. 1998. Ecological considerations of mycorrhizal symbioses. In: Lynch JP, Deikman J, eds. *Phosphorus in plant biology: regulatory roles in molecular, cellular, organismic and ecosystem processes*. American Society of Plant Physiologists Series 19. Rockville, MD, USA: American Society of Plant Physiologists, 17–25.
- Koide RT, Li M. 1989. Appropriate controls for vesicular–arbuscular mycorrhizal research. *New Phytologist* 111: 35–44.
- Koide RT, Lu X. 1992. Mycorrhizal infection of wild oats: maternal effects on offspring growth and reproduction. *Oecologia* 90: 218–225.
- Koide RT, Mooney HA. 1987. Spatial variation in inoculum potential of vesicular–arbuscular mycorrhizal fungi caused by formation of gopher mounds. *New Phytologist* 107: 173–182.
- Koide RT, Li M, Lewis JD, Irby C. 1988. Role of mycorrhizal infection in the growth and reproduction of wild vs cultivated plants. I. Wild vs cultivated oats. *Oecologia* 77: 537–543.
- Lau T-C, Stephenson AG. 1993. Effects of soil nitrogen on pollen production, pollen grain size, and pollen performance in *Cucurbita pepo* (Cucurbitaceae). *American Journal of Botany* 80: 763–768.
- Lau T-C, Stephenson AG. 1994. Effects of soil phosphorus on pollen production, pollen size, pollen phosphorus content, and the ability to sire seeds in *Cucurbita pepo* (Cucurbitaceae). *Sexual Plant Reproduction* 7: 215–220.
- Lau T-C, Lu X, Koide RT, Stephenson AG. 1995. Effects of soil fertility and mycorrhizal infection on pollen production and pollen grain size of *Cucurbita pepo* (Cucurbitaceae). *Plant, Cell & Environment* 18: 169–178.
- Law R. 1988. Evolution in a mutualistic environment. In: Boucher DH, ed. *The biology of mutualism, ecology and evolution*. Oxford, UK: Oxford University Press, 145–170.
- Lee TD, Bazzaz FA. 1982. Regulation of fruit maturation pattern in annual legume, *Cassia fasciculata*. *Ecology* 63: 1374–1388.
- Lewis JD, Koide RT. 1990. Phosphorus supply, mycorrhizal infection and plant offspring vigour. *Functional Ecology* 4: 695–702.
- Lloyd DG. 1980. Sexual strategies in plants. I. An hypothesis of maternal investment during one reproductive session. *New Phytologist* 86: 69–79.
- Lu X. 1993. *Maternal effects of mycorrhizal infection on offspring growth and reproduction*. PhD thesis, The Pennsylvania State University, University Park, PA, USA.
- Lu X, Koide RT. 1991. *Avena fatua* L. seed and seedling nutrient dynamics as influenced by mycorrhizal infection of the maternal generation. *Plant, Cell & Environment* 14: 931–939.
- Lu X, Koide RT. 1994. The effects of mycorrhizal infection on components of plant growth and reproduction. *New Phytologist* 128: 211–218.
- Machlis L, Torrey JG. 1956. *Plants in action*. San Francisco, CA, USA: Freeman.
- Minitab. 1997. *MINITAB reference manual*. Lebanon, PA, USA: Sowers Printing Company.
- Patton DC, Ford HA. 1983. The influence of plant characters and honeyeater size on levels of pollination in Australian plants. In: Jones CE, Little RJ, eds. *Handbook of experimental pollination biology*. New York, NY, USA: van Nostrand Reinhold, 235–248.
- Pendleton RL. 2000. Pre-inoculation by an arbuscular mycorrhizal fungus enhances male reproductive output of *Cucurbita foetidissima*. *International Journal of Plant Science* 161: 683–689.
- Philip LJ, Posluszny U, Klironomos JN. 2001. The influence of mycorrhizal colonization on the vegetative growth and sexual reproductive potential of *Lythrum salicaria* L. *Canadian Journal of Botany* 79: 381–388.
- Poulton JL. 2000. Effects of mycorrhizal infection and soil phosphorus availability on the male function of reproduction in *Lycopersicon esculentum* Mill. PhD thesis, The Pennsylvania State University, University Park, PA, USA.
- Poulton JL, Bryla DR, Koide RT, Stephenson AG. 1998. Effects of mycorrhizal infection and soil phosphorus availability on leaf area measurements in tomato. In: Lynch JP, Deikman J, eds. *Phosphorus in plant biology: regulatory roles in molecular, cellular, organismic and ecosystem processes*. American Society of Plant Physiologists Series 19. Rockville, MD, USA: American Society of Plant Physiologists, 314–315.
- Poulton JL, Koide RT, Stephenson AG. 2001a. Effects of mycorrhizal infection and soil phosphorus availability on in vitro and in vivo pollen performance in *Lycopersicon esculentum* (Solanaceae). *American Journal of Botany* 88: 1786–1793.
- Poulton JL, Koide RT, Stephenson AG. 2001b. Effects of mycorrhizal infection, soil phosphorus availability and fruit production on the male function in two cultivars of *Lycopersicon esculentum*. *Plant, Cell & Environment* 24: 841–849.
- Quesada MR, Bollman K, Stephenson AG. 1995. Leaf damage decreases pollen production and hinders pollen performance in *Cucurbita texana*. *Ecology* 76: 437–443.
- Roach DA, Wulff RD. 1987. Maternal effects in plants. *Annual Review of Ecology and Systematics* 18: 209–236.
- Rovira AD, Bowen GD. 1966. The effects of microorganisms on plant growth. Detoxification of heat-sterilized soils by fungi and bacteria. *Plant Soil* 25: 129–141.
- Schaffer WH, Schaffer MV. 1979. The adaptive significance of variations in reproductive habit in the Agavaceae II. Pollinator foraging behavior and selection for increased reproductive expenditure. *Ecology* 60: 1051–1069.
- Schemske DW. 1980a. Evolution of floral display in the orchid *Brassavola nodosa*. *Evolution* 34: 489–493.
- Schemske DW. 1980b. Floral ecology and hummingbird pollination of *Combretum farinosum* in Costa Rica. *Biotropica* 12: 169–181.
- Schenck NC, Smith GS. 1982. Responses of six species of vesicular–arbuscular mycorrhizal fungi and their effects on soybean at four soil temperatures. *New Phytologist* 92: 193–201.
- Schoen DJ, Steward SC. 1986. Variation in male reproductive investment and male reproductive success in white spruce. *Evolution* 40: 1109–1121.
- Shumway DL, Koide RT. 1994a. Reproductive responses to mycorrhizal colonization of *Abutilon theophrasti* Medic. plants grown for two generations in the field. *New Phytologist* 128: 219–224.
- Shumway DL, Koide RT. 1994b. Within-season variability in mycorrhizal benefit to reproduction in *Abutilon theophrasti* Medic. *Plant, Cell & Environment* 17: 821–827.
- Smith SE, Read DJ. 1997. *Mycorrhiza symbiosis*, 2nd edn. San Diego, CA, USA: Academic Press.
- Solbrig OT. 1981. Studies on the population biology of the genus *Viola* II. The effect of plant size on fitness in *Viola sororia*. *Evolution* 35: 1080–1094.
- Stanley MR, Koide RT, Shumway DL. 1993. Mycorrhizal symbiosis increases growth, reproduction and recruitment of *Abutilon theophrasti* Medic. in the field. *Oecologia* 94: 30–35.
- Stanley RG, Linskens HF. 1974. *Pollen: biology, biochemistry, and management*. Berlin, Germany: Springer.
- Stephenson AG. 1979. An evolutionary examination of the floral display of *Catalpa speciosa* (Bignoniaceae). *Evolution* 33: 1200–1209.

- Stephenson AG. 1992. The regulation of maternal investment in plants. In: Marshall C, Grace J, eds. *Fruit and seed production*. Cambridge, UK: Cambridge University Press, 151–172.
- Stephenson AG, Lau T-C, Quesada M, Winsor JA. 1992. Factors that influence pollen performance. In: Wyatt R, ed. *Ecology and evolution of plant reproduction*. New York, NY, USA: Chapman & Hall.
- Stephenson AG, Erickson CW, Lau T-C, Quesada MR, Winsor JA. 1994. Effects of growing conditions on the male gametophyte. In: Stephenson AG, Kao T-H, eds. *Pollen–pistil interactions and pollen tube growth*. American Society of Plant Physiologists. Series 12. Rockville, MD, USA: American Society of Plant Physiologists, 220–229.
- Stephenson AG, Poulton JL, Lau T-C, Koide RT. 1998. Effects of soil phosphorus level and mycorrhizal infection on the male function of plants. In: Lynch JP, Deikman J, eds. *Phosphorus in plant biology: regulatory roles in molecular, cellular, organismic and ecosystem processes*. American Society of Plant Physiologists Series 19. Rockville, MD, USA: American Society of Plant Physiologists, 52–67.
- Subramanian KS, Charest C. 1997. Nutritional, growth, and reproductive responses of maize (*Zea mays* L.) to arbuscular mycorrhizal inoculation during and after drought stress at tasseling. *Mycorrhiza* 7: 25–32.
- Vasek FC, Weng V, Beaver RJ, Huszar CK. 1987. Effects of mineral nutrition on components of reproduction in *Clarkia unguiculata*. *Aliso* 11: 599–618.
- Vejsadova D, Siblikova D, Gryndler M, Simon T, Miksik I. 1993. Influence of inoculation with *Bradyrhizobium japonicum* and *Glomus claroideum* on seed yield of soybean under greenhouse and field conditions. *Journal of Plant Nutrition* 16: 619–629.
- Watanabe FS, Olsen SR. 1965. Test of an ascorbic acid method for determining phosphorus in water and NaHCO₃ extracts from soil. *Soil Science Society Proceedings* 29: 677–678.
- Willson MF, Miller LJ, Rathcke BJ. 1979. Floral displays in *Phlox* and *Geranium*: adaptive aspects. *Evolution* 33: 52–63.
- Young HJ, Stanton ML. 1990. Influence of environmental quality on pollen competitive ability in wild radish. *Science* 248: 1631–1633.



About *New Phytologist*

- *New Phytologist* is owned by a non-profit-making **charitable trust** dedicated to the promotion of plant science. Regular papers, Letters, Research reviews, Rapid reports and Methods papers are encouraged. Complete information is available at www.newphytologist.com
- All the following are **free** – essential colour costs, 100 offprints for each article, online summaries and ToC alerts (go to the website and click on Synergy)
- You can take out a **personal subscription** to the journal for a fraction of the institutional price. Rates start at £83 in Europe/\$133 in the USA & Canada for the online edition (go to the website and click on Subscriptions)
- If you have any questions, do get in touch with Central Office (newphytol@lancaster.ac.uk; tel +44 1524 594691) or, for a local contact in North America, the USA Office (newphytol@ornl.gov; tel 865 576 5251)